arXiv:math/0010190v1 [math.QA] 19 Oct 2000

On quantization of r-matrices for Belavin-Drinfeld Triples

A.P. Isaev* and O.V.Ogievetsky†

* *Bogoliubov Laboratory of Theoretical Physics, JINR, 141980 Dubna,*
Moscow region, Russia

† *Center of Theoretical Physics, Luminy, 13288 Marseille, France*
and
P. N. Lebedev Physical Institute, Theoretical Department, Leninsky pr. 53,
117924 Moscow, Russia

Abstract

We suggest a formula for quantum universal R-matrices corresponding to quasitriangular classical r-matrices classified by Belavin and Drinfeld for all simple Lie algebras. The R-matrices are obtained by twisting the standard universal R-matrix.

1. Classical quasitriangular r-matrices for semisimple Lie algebras are classified by Belavin-Drinfeld triples [1]. The Belavin-Drinfeld triple $(\Gamma_1,\,\Gamma_2,\,\tau)$ for a simple Lie algebra $\mathbf{g}=\mathbf{g}^+\oplus\mathbf{h}\oplus\mathbf{g}^-$ consists of the following data: Γ_1, Γ_2 are subsets of the set Γ of simple roots of the algebra $\mathbf{g}$ and τ is a one-to-one mapping: $\Gamma_1\rightarrow\Gamma_2$ such that $<\tau(\alpha),\,\tau(\beta)>=<\alpha,\,\beta>$ and $\tau^k(\alpha)\neq\alpha$ for any $\alpha,\beta\in\Gamma_1$ and any natural k. The corresponding quantum R-matrices should have the form

$$R_{12}=F_{21}\,\mathcal{R}_{12}\,F_{12}^{-1}\;, \tag{1}$$

where $\mathcal{R}$ is the standard universal Drinfeld-Jimbo R-matrix for the Lie algebra $\mathbf{g}$. The twisting operator satisfies the cocycle equation

$$F_{12}\,(\Delta\otimes id)\,F=F_{23}\,(id\otimes\Delta)\,F\;. \tag{2}$$

Therefore the problem of quantization is reduced to the problem of finding the twisting operator F_{12} for each Belavin-Drinfeld triple. In the present paper we suggest a formula for the twisting operator F_{12}. We present the twisting operator in a factorized form

$$F_{12}=F_{12}^{(N)}\cdot F_{12}^{(N-1)}\cdots F_{12}^{(2)}\cdot F_{12}^{(1)}\cdot K\;, \tag{3}$$

where the factors $F^{(k)}$ are special canonical elements defined by the powers of the one-to-one map τ; the operator K belongs to $q^{\mathbf{h}\otimes\mathbf{h}}$. A different formula for the operator F_{12} was given in [2]. We shall say several words about the differences at the end of the present paper.

The plan of our paper is as follows.

Our approach heavily uses the modified Cartan-Weyl basis for $U_q(\mathbf{g})$. The definition of the modified simple root generators is contained in Section 2. In Section 3 we give an interpretation of Belavin-Drinfeld triples in terms of the modified basis. In Section 4 a modified Cartan-Weyl basis is introduced. The twisting operator F_{12} is constructed in Section 5. Finally, in Section 6 several examples are presented.

Everywhere below we assume the deformation parameter q to be generic (not a root of unity).

2. *Modified basis for quantum universal enveloping algebras.*

Consider a quantum universal enveloping algebra $U_q(\mathbf{g})$ with relations (see e.g. [3])

$$[h_i,\, h_j] = 0 \ , \quad [h_i,\, e_j] = a_{ij}\, e_j \ , \quad [h_i,\, f_j] = -a_{ij}\, f_j \ ,$$

$$[e_i,\, f_j] = \delta_{ij}\, \frac{K_i - K_i^{-1}}{q^{d_i} - q^{-d_i}} \ , \tag{4}$$

and Serre relations

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \left[\begin{array}{c} 1-a_{ij} \\ k \end{array} \right]_{q^{d_i}} (e_i)^k\, e_j\, (e_i)^{1-a_{ij}-k} = 0 \ , \tag{5}$$

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \left[\begin{array}{c} 1-a_{ij} \\ k \end{array} \right]_{q^{d_i}} (f_i)^k\, f_j\, (f_i)^{1-a_{ij}-k} = 0 \ , \tag{6}$$

where

$$\left[\begin{array}{c} n \\ k \end{array} \right]_q = \frac{[n]_q!}{[k]_q!\,[n-k]_q!} \ , \quad [k]_q = \frac{q^k - q^{-k}}{q - q^{-1}} \ ,$$

a_{ij} is the Cartan matrix for $\mathbf{g}$, $K_i = q^{d_i h_i}$ and d_i are smallest positive integers (from the set $1, 2, 3$) such that $d_i a_{ij} = a^{(s)}_{ij}$ is symmetric matrix. The algebra $U_q(\mathbf{g})$ is a Hopf algebra with the comultiplication

$$\begin{array}{c} \Delta(h_i) = h_i \otimes 1 + 1 \otimes h_i \ , \\ \Delta(e_i) = e_i \otimes K_i + 1 \otimes e_i \ , \quad \Delta(f_i) = f_i \otimes 1 + K_i^{-1} \otimes f_i \ . \end{array} \tag{7}$$

The antipode and the counit are

$$S(h_i) = -h_i \ , \quad S(e_i) = -e_i\, K_i^{-1} \ , \quad S(f_i) = -K_i\, f_i \ ,$$

$$\epsilon(h_i) = \epsilon(e_i) = \epsilon(f_i) = 0 \ .$$

Any operator $K \in q^{\mathbf{h} \otimes \mathbf{h}}$,

$$K = q^{\left(\sum_{ij} b_{ij}\, h_i \otimes h_j\right)} \ , \tag{8}$$

for arbitrary numerical matrix b_{ij}, obviously satisfies the cocycle equation (2),

$$K_{12}\, (\Delta \otimes id)\, K = K_{23}\, (id \otimes \Delta)\, K \ . \tag{9}$$

Therefore one can twist the comultiplication by K:

$$\widetilde{\Delta}(a) := K\,\Delta(a)\,K^{-1}\ . \tag{10}$$

We change the basis in the algebra $U_q(\mathbf{g})$ by introducing new generators

$$E_i = X_i\,e_i\ , \quad F_i = f_i\,Y_i\ , \tag{11}$$

where $X_i = \exp(\sum_j x_{ij}\,h_j)$, $Y_i = \exp(\sum_j y_{ij}\,h_j)$ and x_{ij}, y_{ij} are some numerical matrices. We require that the comultiplication (10) for the new generators (11) has the following form:

$$\begin{aligned} \widetilde{\Delta}(E_i) &= K\,\Delta(E_i)\,K^{-1} = E_i \otimes R_i^+ + 1 \otimes E_i\ , \\ \widetilde{\Delta}(F_i) &= K\,\Delta(F_i)\,K^{-1} = F_i \otimes 1 + R_i^- \otimes F_i\ . \end{aligned} \tag{12}$$

Equations (12) relate operators X_i, Y_i and K.

A comparison of (7) and (12) gives

$$X_i = q^{-\sum_{mn} h_m b_{mn} a_{ni}} \equiv q^{-(h\,b\,a)_i}\ , \quad Y_i = q^{\sum_{mn} h_m b_{nm} a_{ni}} \equiv q^{(h\,\overline{b}\,a)_i}\ ,$$

and

$$R_i^{\pm} = X_i\,K_i^{\pm 1}\,Y_i = K_i^{\pm 1}\,q^{-(h(b-\overline{b})a)_i}\ , \quad R_i^+ = K_i^2\,R_i^-\ , \tag{13}$$

where $\overline{b}_{mn} = b_{nm}$ is the transposed matrix.

The relations (4) and Serre relations (5), (6) for the quantum algebra $U_q(g)$ in terms of the new generators (11) take the form

$$[E_i,\,F_j] = \delta_{ij}\frac{R_i^+ - R_i^-}{q^{d_i} - q^{-d_i}}\ , \tag{14}$$

$$R_i^{\pm}\,E_j = q^{\pm a_{ij}^{(s)} + A_{ij}}\,E_j\,R_i^{\pm}\ , \quad R_i^{\pm}\,F_j = q^{\mp a_{ij}^{(s)} - A_{ij}}\,F_j\,R_i^{\pm}\ , \tag{15}$$

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_{q^{d_i}} q^{-kA_{ij}}\,(E_i)^k\,E_j\,(E_i)^{1-a_{ij}-k} = 0\ , \tag{16}$$

$$\sum_{k=0}^{1-a_{ij}} (-1)^k \begin{bmatrix} 1-a_{ij} \\ k \end{bmatrix}_{q^{d_i}} q^{kA_{ij}}\,(F_i)^k\,F_j\,(F_i)^{1-a_{ij}-k} = 0\ , \tag{17}$$

with a skewsymmetric matrix $A_{ij} = (\overline{a}(b-\overline{b})a)_{ij}$.

In the sequel we shall use q-commutators: $[A,B]_\mu := A\,B - \mu\,B\,A$. Relations (16), (17) can be conveniently rewritten in terms of q-commutators. For example, for $a_{ij}=0$ the relations $[e_i,e_j]=0=[f_i,f_j]$ are rewritten as

$$[E_i,E_j]_{q^{A_{ij}}} = 0\ , \quad [F_i,F_j]_{q^{-A_{ij}}} = 0\ ,$$

while for $a_{ij}=-1$ we have

$$\begin{aligned} &[[E_i,E_j]_\mu,E_i]_\nu = 0 = [E_j,[E_i,E_j]_\mu]_\nu\ , \\ &[[F_i,F_j]_\nu,F_i]_\mu = 0 = [F_j,[F_i,F_j]_\nu]_\mu\ . \end{aligned} \tag{18}$$

where $\mu = q^{d_i + A_{i\,j}}$, $\nu = q^{d_i - A_{i\,j}}$.

Remark. The modified basis for multiparametric twistings of $U_q(\mathbf{g})$ has been considered by T.Hodges [4].

3. *Modified basis and Belavin - Drinfeld triples.*

All the data from the Belavin-Drinfeld triple can be conveniently interpreted in terms of the modified basis for a suitable matrix b_{ij}:

Proposition. *Let* Γ *be the set of simple roots of* $\mathbf{g}$, Γ_1 *and* Γ_2 *subsets of* Γ *and* τ *a one-to-one mapping:* $\Gamma_1 \rightarrow \Gamma_2$. *Then the following equations for the matrix* b_{ij}

$$R^+_{\alpha_i} = R^-_{\tau(\alpha_i)}\ \ \forall\ \alpha_i \in \Gamma_1 \tag{19}$$

where $R^\pm_{\alpha_i} \equiv R^\pm_i$, *admit a solution if and only if the triple* $(\Gamma_1, \Gamma_2, \tau)$ *is the Belavin-Drinfeld triple.*

Proof. Assume that a solution of equation (19) exists. We then need to prove that the mapping τ satisfies conditions:

$$\text{1) for any } \alpha \in \Gamma_1 \text{ there is a natural } k \text{ for which } \tau^k(\alpha) \notin \Gamma_1\,, \tag{20}$$

$$\text{2) for any } \alpha,\ \beta \in \Gamma_1\ ,\ \ <\tau(\alpha),\tau(\beta)> = <\alpha,\beta>\ \ . \tag{21}$$

The condition (20) means that τ has no cycles: $\tau^k(\alpha)) \neq \alpha$ for all $\alpha \in \Gamma_1$ and $k > 0$.

Indeed, assume that τ has a cycle, $\tau^k(\alpha) = \alpha$ for some $\alpha \in \Gamma_1$ and a natural k. Take a minimal k with this property.

Then $R^+_{\tau^k(\alpha)} = R^+_\alpha$ and equations (13) imply

$$\begin{aligned} R^+_\alpha = R^-_{\tau(\alpha)} = K^{-2}_{\tau(\alpha)}\, R^+_{\tau(\alpha)} = \ldots = \left(\textstyle\prod_{i=1}^k\, K^{-2}_{\tau^i(\alpha)}\right)\, R^+_{\tau^k(\alpha)} \\ = \left(\textstyle\prod_{i=1}^k\, K^{-2}_{\tau^i(\alpha)}\right)\, R^+_\alpha\ . \end{aligned} \tag{22}$$

Therefore

$$\left(\prod_{i=1}^k K^{-2}_{\tau^i(\alpha)}\right) = 1\ ,$$

which contradicts to the independence of generators in the Cartan subalgebra of $U_q(\mathbf{g})$. Thus, $\tau^k(\alpha)$ never equals α which proves the condition (20).

To prove the condition (21), note that eq. (19) is equivalent to the following condition on the skewsymmetric matrix $A_{mn} = (\overline{a}(b-\overline{b})a)_{mn}$

$$A_{im} + A_{m\tau(i)} + a^{(s)}_{im} + a^{(s)}_{\tau(i)m} = 0\ , \tag{23}$$

where the subscript m runs over all simple roots while i numerates only roots from Γ_1. Eq. (23) is obtained by commuting both sides of eq. (19) with e_m (or f_m). Here it is important that q is not a root of unity.

For indices i, m corresponding to roots $\alpha_i, \alpha_m \in \Gamma_1$, eq. (23) can be rewritten in the following three equivalent forms

$$A_{i\tau(m)} + A_{\tau(m)\tau(i)} + a^{(s)}_{i\tau(m)} + a^{(s)}_{\tau(i)\tau(m)} = 0\ , \tag{24}$$

$$A_{mi} + A_{i\tau(m)} + a^{(s)}_{mi} + a^{(s)}_{\tau(m)i} = 0\ , \tag{25}$$

$$A_{m\tau(i)} + A_{\tau(i)\tau(m)} + a^{(s)}_{m\tau(i)} + a^{(s)}_{\tau(m)\tau(i)} = 0\ , \tag{26}$$

The combinations (23) + (25) and (24) + (26) of the equations are, respectively,

$$2a^{(s)}_{im} = -a^{(s)}_{\tau(m)i} - a^{(s)}_{\tau(i)m} - A_{m\tau(i)} - A_{i\tau(m)}\ , \tag{27}$$

$$2a^{(s)}_{\tau(i)\tau(m)} = -a^{(s)}_{i\tau(m)} - a^{(s)}_{m\tau(i)} - A_{m\tau(i)} - A_{i\tau(m)}\ . \tag{28}$$

Therefore,

$$a^{(s)}_{im} = a^{(s)}_{\tau(i)\tau(m)}\ , \tag{29}$$

which is equivalent to the second condition $< \tau(\alpha_i), \tau(\alpha_m) >=< \alpha_i, \alpha_m >$ for the Belavin-Drinfeld triple. •

Remark 1. The difference of eqs. (24) and (25) gives the following relation on the matrix A_{ij}

$$A_{im} - A_{\tau(i)\tau(m)} = a^{(s)}_{mi} + a^{(s)}_{\tau(m)i} - a^{(s)}_{i\tau(m)} - a^{(s)}_{\tau(i)\tau(m)} = 0 \ . \tag{30}$$

This shows that the map τ does not change the modified basis.

Remark 2. Consider two sequences of sets

$$\begin{aligned} &\Gamma_1 = \Gamma_1^{(0)} \supset \Gamma_1^{(1)} \supset \Gamma_1^{(2)} \ldots \supset \Gamma_1^{(N)} \supset \Gamma_1^{(N+1)} = \emptyset \ , \\ &\Gamma_2 = \Gamma_2^{(0)} \supset \Gamma_2^{(1)} \supset \Gamma_2^{(2)} \ldots \supset \Gamma_2^{(N)} \ , \end{aligned} \tag{31}$$

defined by

$$\Gamma_1^{(k+1)} = \Gamma_1^{(k)} \cap \Gamma_2^{(k)} \ , \quad \Gamma_1^{(k)} \xrightarrow{\tau} \Gamma_2^{(k)} \ .$$

We assume that the set $\Gamma_1^{(N)}$ is not empty. The number N is called the degree of the triple $(\Gamma_1,\, \Gamma_2,\, \tau)$.

Introduce a set $\widetilde{\Gamma}_1^{(k)} = \tau^{-k-1}(\Gamma_2^{(k)}) \in \Gamma_1$. Then the mapping τ^k: $\widetilde{\Gamma}_1^{(k-1)} \xrightarrow{\tau^k} \Gamma_2^{(k-1)} \neq \emptyset$ also defines a Belavin-Drinfeld triple

$$(\widetilde{\Gamma}_1^{(k-1)},\, \Gamma_2^{(k-1)}, \tau^k) \ . \tag{32}$$

4. *Modified Cartan-Weyl basis and normal order of roots.*

Let Δ_+ be the system of all positive roots of $\mathbf{g}$ with respect to Γ. A construction of Cartan-Weyl basis in terms of the modified generators E_i and F_i is analogous to the usual procedure for $U_q(\mathbf{g})$ (see [5]).

Recall the notion of a normal (convex) order in Δ_+: the set Δ_+ is ordered normally if any root γ which is a sum of roots α and β is placed between α and β.

We write $\alpha < \beta$ if the root α is located to the left of the root β. For $\alpha < \beta$, the interval between roots α and β is denoted by $\{\alpha, \beta\}$.

Given a normal order in Δ_+, the modified Cartan-Weyl basis is constructed by the following inductive procedure. The generators for the simple roots are already defined. For a composite root γ, take a minimal interval $\{\alpha, \beta\}$, $\alpha < \beta$, with $\gamma = \alpha + \beta$ ("minimal" means that there is no subinterval $\{\widetilde{\alpha}, \widetilde{\beta}\} \subset \{\alpha, \beta\}$ for which $\gamma = \widetilde{\alpha} + \widetilde{\beta}$). Assume that generators E_α, E_β, F_α and F_β were defined at previous steps. Then generators E_γ and F_γ are defined by

$$E_\gamma = [E_\alpha,\, E_\beta]_\mu = E_\alpha\, E_\beta - \mu E_\beta\, E_\alpha\ , \tag{33}$$

$$F_\gamma = [F_\alpha,\, F_\beta]_\nu = F_\alpha\, F_\beta - \nu F_\beta\, F_\alpha\ . \tag{34}$$

where

$$\mu = q^{-<\alpha,\beta>+<\alpha,\, A\,\beta>}\ , \quad \nu = q^{-<\alpha,\beta>-<\alpha,\, A\,\beta>}$$

and A is the operator with the matrix A_{ij}:

$$< \alpha_i,\, A\, \alpha_j >= A_{ij}\ .$$

If there are several possible minimal intervals $\{\alpha, \beta\}$ for which $\gamma = \alpha + \beta$, the definitions (33)-(34) give proportional results.

Note. For the case $A_{ij} = 0$ the definition (33)-(34) of composite roots does not coincide with the definition in [5] since we use the comultiplication (7) which is different from the comultiplication in [5].

5. *Twisting operators F_{12} for Belavin-Drinfeld triples.*

For a given simple Lie algebra $\mathbf{g}$ fix a normal order in Δ_+.

We need the expression for the inverse of the universal R- matrix for the algebra $U_q(\mathbf{g})$:

$$\mathcal{R}^{-1} = \vec{\prod}_{\beta\in\Delta_+}\ \exp_{q_\beta}\left(-\lambda\, a_\beta\, (e_\beta \otimes f_\beta)\right) \cdot K^{(0)}\ , \tag{35}$$

where $q_\alpha = q^{<\alpha,\alpha>}$, $\lambda = q - q^{-1}$ and $K^{(0)} \in q^{\mathbf{h}\otimes\mathbf{h}}$. The product in eq. (35) is the ordered product corresponding to the chosen normal order of roots. For precise values of the constants a_β see [6], [5]. The function $\exp_q$ is the standard q-exponent,

$$\exp_q(u) = \prod_{n=0}^{\infty}(1 + (q-1)uq^n)^{-1} = \sum_{k=0}^{\infty} \frac{u^k}{k_q!}\ , \quad k_q = \frac{q^k - 1}{q - 1}\ . \tag{36}$$

Let $(\Gamma_1,\, \Gamma_2,\, \tau)$ be a Belavin-Drinfeld triple of degree N. Define elements $F^{(k)}$ by

$$F_{12}^{(k)} = \vec{\prod}_{\beta\in\Delta_+^{(k)}} \ \exp_{q_\beta} \Big(-\lambda\, a_\beta \, (E_\beta \otimes F_{\tau^k(\beta)}) \Big) \ , \tag{37}$$

where in the ordered product we keep terms corresponding to only those roots β for which $\tau^k(\beta)$ is defined (that is, the element e_β belongs to the subalgebra with generators from the subset $\widetilde{\Gamma}_1^{(k)}$ defined in (32)). This is reflected in the notation $\beta \in \Delta_+^{(k)}$.

The expression (37) can be given a form

$$F_{12}^{(k)} = (1 \otimes T^k) \Big(K\, \mathcal{R}^{-1}\, (K^{(0)})^{-1}\, K^{-1} \Big) \ , \tag{38}$$

where the operator T on the elements F_β is defined by $T(F_\beta) = F_{\tau(\beta)}$ wherever $\tau(\beta)$ is defined; $T(F_\beta) = 0$ otherwise. The operator K corresponds to the solution of eqs. (19) for the given Belavin-Drinfeld triple.

Theorem. *For the quantum algebra $U_q(\mathbf{g})$ and the Belavin-Drinfeld triple $(\Gamma_1,\, \Gamma_2,\, \tau)$ of degree N the universal twisting element F_{12} is*

$$F_{12} = F_{12}^{(N)} \cdot F_{12}^{(N-1)} \cdots F_{12}^{(2)} \cdot F_{12}^{(1)} \cdot K \equiv \widetilde{F}_{12} \cdot K \tag{39}$$

with the factors $F_{12}^{(k)}$ defined in (37).

We sketch the proof shortly. It is based on explicit formulas for the coproduct of elements $F_{12}^{(k)}$:

$$(\widetilde{\Delta} \otimes id) F_{12}^{(k)} = F_{23}^{(k)}\, (K_{23}^{(k)})^{-1}\, F_{13}^{(k)}\, K_{23}^{(k)} \ , \tag{40}$$

$$(id \otimes \widetilde{\Delta}) F_{12}^{(k)} = F_{12}^{(k)}\, (K_{12}^{(k)})^{-1}\, F_{13}^{(k)}\, K_{12}^{(k)} \ , \tag{41}$$

for some elements $K^{(k)} \in q^{\mathbf{h}\otimes\mathbf{h}}$. The comultiplication $\widetilde{\Delta}$ is twisted as in (10).

Next, one can verify the following identities

$$\widetilde{F}_{23}^{(k)}\, \widetilde{F}_{13}^{(k+m)}\, \widetilde{F}_{12}^{(m)} = \widetilde{F}_{12}^{(m)}\, \widetilde{F}_{13}^{(k+m)}\, \widetilde{F}_{23}^{(k)} \ , \tag{42}$$

where $\widetilde{F}^{(k)} = F^{(k)} \cdot (K^{(k)})^{-1}$.

With the help of (40) - (42) it is straightforward to check the cocycle condition (2).

Remark 1. Another expression for the twisting element F was suggested in [2]. The expression in [2] has a factorised form as well. However, the factors $F^{(i)}$ are different; one of the differences is that each factor in [2] contains terms from $q^{\mathbf{h}\otimes\mathbf{h}}$. In our expression (39) all terms from $q^{\mathbf{h}\otimes\mathbf{h}}$ are collected; the price is the appearance of the modified basis.

Remark 2. The element F in (39) satisfies the following analogue of the linear ABRR equation [7]:

$$(1\otimes T)(F_{12}\mathcal{R}^{-1}(K^{(0)})^{-1}K^{-1}) = F_{12}K^{-1}\ . \tag{43}$$

6. *Examples.*

i) $U_q(sl(3))$ **case** (see [4]).
Here we have only one nontrivial Belavin-Drinfeld triple:



Fig.1

This Cremmer-Gervais type triple has degree 1 and the basic relations (19) which define this triple are reduced to one equation $R_1^+ = R_2^-$. The antisymmetric matrix A_{ij} is

$$A_{ij} = \delta_{i,j+1} - \delta_{j,i+1}\ \ , \tag{44}$$

with $1\le i,j\le 2$. The corresponding universal twisting element (39) has the form

$$F_{12} = F_{12}^{(1)}\cdot K = \exp_{q^2}(-\lambda\, E_1\otimes F_2)\cdot K\ . \tag{45}$$

ii) Cremmer-Gervais $U_q(sl(4))$ **case.**
For this case the triple is given by the following diagram



Fig.2

It has degree 2. The basic relations (19) which define this triple are $R_1^+ = R_2^-$, $R_2^+ = R_3^-$. The matrix A_{ij} is given by (44), now with $1 \leq i, j \leq 2$. The corresponding universal twisting element (39) has the form

$$F_{12} = F_{12}^{(2)} \cdot F_{12}^{(1)} \cdot K \ , \tag{46}$$

where

$$F_{12}^{(2)} = \exp_{q^2}(-\lambda \, E_1 \otimes F_3) \ , \tag{47}$$

$$F_{12}^{(1)} = \exp_{q^2}(-\lambda \, E_1 \otimes F_2) \, \exp_{q^2}(q^{-1}\lambda \, [E_{12}] \otimes [F_{23}]_{q^2}) \, \exp_{q^2}(-\lambda E_2 \otimes F_3) \ . \tag{48}$$

Here $[E_{12}] = E_1 E_2 - E_2 E_1$ and $[F_{23}]_{q^2} = F_2 F_3 - q^2 F_3 F_2$.

Remark. One can directly check that (45), (46) obeys the cocycle conditions (2). For (45) this check requires only the basic equation for the q-exponent,

$$\exp_q(y) \, \exp_q(x) = \exp_q(x + y) \quad \text{if} \quad x \, y = q \, y \, x \ . \tag{49}$$

For (46) one needs two more quantum identities. The first one is the famous pentagon identity (see e.g. [8] and references therein)

$$\exp_q(u) \, \exp_q(v) = \exp_q(v) \, \exp_q([u, v]) \, \exp_q(u) \ , \tag{50}$$

where the operators u and v satisfy the commutation (Serre) relations

$$u \, [u, v] = q \, [u, v] \, u \ , \quad v \, [u, v] = q^{-1} \, [u, v] \, v \ .$$

The second identity is

$$\exp_{q^2}(E) \, \exp_{q^2}(-R^+) \, \exp_{q^2}(F) = \exp_{q^2}(F) \, \exp_{q^2}(-R^-) \, \exp_{q^2}(E) \ , \tag{51}$$

where E, F and $R^\pm$ generate the algebra

$$[E, F] = (R^+ - R^-) \ , \quad [R^+, \, R^-] = 0 \ ,$$
$$R^\pm \, E = q^{\pm 2} \, E \, R^\pm \ , \quad R^\pm \, F = q^{\mp 2} \, F \, R^\pm \ .$$

Acknowledgments: We are indebted to V. Fock, S. Khoroshkin, S. Pakuliak, V. Tolstoy and especially P.N. Pyatov for valuable discussions and comments. One of the authors (API) thanks the Centre de Physique Theorique in Luminy (Marseille University) for the hospitality during his visit in April 1999 where the considerable part of the results of this paper has been obtained. This work was supported in part by the CNRS grant PICS No. 608 and the RFBR grant No. 98-01-2033.

References

[1] A. A. Belavin and V. G. Drinfeld, *Triangle Equations and Simple Lie Algebras*, Sov. Sci. Rev. **C** 4 (1984) 93-166.

[2] P. Etingof, T. Schedler and O. Schiffmann, *Explicit quantization for dynamical r-matrices for finite dimensional semisimple Lie algebras*, (1999), math.QA/9912009.

[3] V.Chari and A.Pressley, *A Guide to Quantum Groups*, Cambridge University Press (1994).

[4] T.Hodges, *Nonstandard quantum groups associated to certain Belavin-Drinfeld triples*, (1996), q-alg/9609029.

[5] S.M.Khoroshkin and V.N.Tolstoy, *Universal R-matrix for Quantized (Super)Algebras* Coom.Math.Phys. **141** (1991) 599; *The uniqueness theorem for the universal R-matrix*, Lett.Math.Phys. **24** (1992) 231; *Twisting of quantum (super)algebras. Connection of Drinfeld's and Cartan-Weyl realizations for quantum affine algebras*, (1994), hep-th/9404036.

[6] M. Rosso, *An analogue of PBW theorem and the universal R- matrix for $U_h(sl(n+1))$*, Commun. Math. Phys. **124** (1989) 307-318.

[7] D. Arnaudon, E. Buffenoir, E. Ragoucy and Ph. Roche, *Universal Solution of Quantum Dynamical Yang-Baxter Equation* Lett. Math. Phys. **44** (1998) 201-214; q-alg/9712037.

[8] L.D.Faddeev and R.M.Kashaev, *Quantum dilogarithm*, Mod.Phys.Lett A, **9** (1994) 427, hep-th/9310070; R.M.Kashaev, *Heisenberg Double and Pentagon Relation*, q-alg/9503005.